

Dr Oliver Mathematics
Advance Level Mathematics
Mechanics 1: Calculator
1 hour 30 minutes

The total number of marks available is 75.

You must write down all the stages in your working.

1. Two particles, P and Q , have masses $3m$ and m respectively. They are moving in opposite directions towards each other along the same straight line on a smooth horizontal plane and collide directly. The speeds of P and Q immediately before the collision are $2u$ and $4u$ respectively. The magnitude of the impulse received by each particle in the collision is $\frac{21mu}{4}$.
 - (a) Find the speed of P after the collision. (3)
 - (b) Find the speed of Q after the collision. (3)
2. A particle of mass 2 kg lies on a rough plane. The plane is inclined to the horizontal at 30° . The coefficient of friction between the particle and the plane is $\frac{1}{4}$. The particle is held in equilibrium by a force of magnitude P newtons. The force makes an angle of 20° with the horizontal and acts in a vertical plane containing a line of greatest slope of the plane, as shown in Figure 1. (10)




Figure 1: a particle of mass 2 kg lies on a rough plane

Find the least possible value of P .

3. A wooden beam AB , of mass 150 kg and length 9 m, rests in a horizontal position supported by two vertical ropes. The ropes are attached to the beam at C and D , where $AC = 1.5$ m and $BD = 3.5$ m. A gymnast of mass 60 kg stands on the beam at the point P , where $AP = 3$ m, as shown in Figure 2. The beam remains horizontal and in equilibrium.




Figure 2: a gymnast of mass 60 kg stands on the beam

By modelling the gymnast as a particle, the beam as a uniform rod and the ropes as light inextensible strings,

- (a) find the tension in the rope attached to the beam at C . (3)

The gymnast at P remains on the beam at P and another gymnast, who is also modelled as a particle, stands on the beam at B . The beam remains horizontal and in equilibrium. The mass of the gymnast at B is the largest possible for which the beam remains horizontal and in equilibrium.

- (b) Find the tension in the rope attached to the beam at D . (4)

4. A ball of mass 0.2 kg is projected vertically downwards with speed U ms^{-1} from a point A which is 2.5 m above horizontal ground. The ball hits the ground. Immediately after hitting the ground, the ball rebounds vertically with a speed of 10 ms^{-1} . The ball receives an impulse of magnitude 7 Ns in its impact with the ground. By modelling the ball as a particle and ignoring air resistance, find

- (a) the value of U . (6)

After hitting the ground, the ball moves vertically upwards and passes through a point B which is 1 m above the ground.

- (b) Find the time between the instant when the ball hits the ground and the instant when the ball first passes through B . (4)

- (c) Sketch a velocity-time graph for the motion of the ball from when it was projected from A to when it first passes through B . (You need not make any further calculations to draw this sketch.) (3)

5. A lift of mass 250 kg is being raised by a vertical cable attached to the top of the lift. A woman of mass 60 kg stands on the horizontal floor inside the lift, as shown in Figure 3.




Figure 3: a lift of mass 250 kg is being raised

The lift ascends vertically with constant acceleration 2 ms^{-2} . There is a constant downwards resistance of magnitude 100 N on the lift. By modelling the woman as a particle,

- (a) find the magnitude of the normal reaction exerted by the floor of the lift on the woman. (3)

The tension in the cable must not exceed 10 000 N for safety reasons, and the maximum upward acceleration of the lift is 3 ms^{-2} . A typical occupant of the lift is modelled as a particle of mass 75 kg and the cable is modelled as a light inextensible string. There is still a constant downwards resistance of magnitude 100 N on the lift.

- (b) Find the maximum number of typical occupants that can be safely carried in the lift when it is ascending with an acceleration of 3 ms^{-2} . (7)

6. (*In this question \mathbf{i} and \mathbf{j} are horizontal unit vectors due east and due north respectively.*) Two forces \mathbf{F}_1 and \mathbf{F}_2 act on a particle P of mass 0.5 kg.

$$\mathbf{F}_1 = (4\mathbf{i} - 6\mathbf{j}) \text{ N and } \mathbf{F}_2 = (p\mathbf{i} + q\mathbf{j}) \text{ N.}$$

Given that the resultant force of \mathbf{F}_1 and \mathbf{F}_2 is in the same direction as $(-2\mathbf{i} - \mathbf{j})$,

- (a) show that $p - 2q = -16$. (5)

Given that $q = 3$,

- (b) find the magnitude of the acceleration of P , (5)
 (c) find the direction of the acceleration of P , giving your answer as a bearing to the nearest degree. (3)

7. A particle P of mass $4m$ is held at rest at the point X on the surface of a rough inclined plane which is fixed to horizontal ground. The point X is a distance h from the bottom of the inclined plane. The plane is inclined to the horizontal at an angle α where $\tan \alpha = \frac{3}{4}$. The coefficient of friction between P and the plane is $\frac{1}{4}$. The particle P is attached to one end of a light inextensible string. The string passes over a small smooth pulley which is fixed at the top of the plane. The other end of the string is attached to a particle Q of mass m which hangs freely at a distance d , where $d > h$, below the pulley, as shown in Figure 4.




Figure 4: a particle P of mass $4m$

The string lies in a vertical plane through a line of greatest slope of the inclined plane. The system is released from rest with the string taut and P moves down the plane. For the motion of the particles before P hits the ground,

- (a) state which of the information given above implies that the magnitudes of the accelerations of the two particles are the same, (1)
- (b) write down an equation of motion for each particle, (5)
- (c) find the acceleration of each particle. (5)

When P hits the ground, it immediately comes to rest. Given that Q comes to instantaneous rest before reaching the pulley,

- (d) show that (5)

$$d > \frac{28h}{25}.$$