

Dr Oliver Mathematics
Mathematics
Integration Part 3
Past Examination Questions

This booklet consists of 67 questions across a variety of examination topics.
The total number of marks available is 451.

1. (a) Express (3)

$$\frac{5x + 3}{(2x - 3)(x + 2)}$$

in partial fractions.

- (b) Hence find the exact value of (5)

$$\int_2^6 \frac{5x + 3}{(2x - 3)(x + 2)} dx,$$

giving your answer as a single logarithm.

2. Use the substitution $x = \sin \theta$ to find the exact value of

$$\int_0^{\frac{1}{2}} \frac{1}{(1 - x^2)^{\frac{3}{2}}} dx.$$

3. Using the substitution $u^2 = 2x - 1$, or otherwise, to find the exact value of

$$\int_1^5 \frac{3x}{\sqrt{2x - 1}} dx.$$

4. The finite region R , shown shaded in the figure, is bounded by the curve $y = xe^x$, the x -axis, and the lines $x = 1$ and $x = 3$.




Figure 1: $y = xe^x$

The region R is rotated through 360 degrees about the x -axis. Use integration by parts to find an exact value for the **volume** of the solid generated.

5. The curve shown in the Figure 2 has parametric equations

$$x = t - 2 \sin t, y = 1 - 2 \cos t.$$




Figure 2: $x = t - 2 \sin t, y = 1 - 2 \cos t$

(a) Show that the curve crosses the x -axis where $t = \frac{\pi}{3}$ and $t = \frac{5\pi}{3}$. (2)

The finite region R is enclosed by the curve and the x -axis.

(b) Show that the area of R is given by the integral (3)

$$\int_{\frac{\pi}{3}}^{\frac{5\pi}{3}} (1 - 2 \cos t)^2 dt.$$

(c) Use this integral to find the exact value of the shaded area. (7)

6. The curve with equation $y = 3 \sin \frac{x}{2}$, $0 \leq x \leq 2\pi$, is shown in Figure 3.




Figure 3: $y = 3 \sin \frac{x}{2}$

The finite region enclosed by the curve and the x -axis is shaded.

- (a) Find, by integration, the area of the shaded region. (3)

This region is rotated through 2π radians about the x -axis.

- (b) Find the volume of the solid generated. (6)

7. Show, by integration, that the exact value of $\int_1^3 (x-1) \ln x \, dx$ is $\frac{3}{2} \ln 3$. (6)

8. The curve with equation $y = \frac{1}{3(1+2x)}$, $x > -\frac{1}{2}$, is shown in Figure 4.




Figure 4: $y = \frac{1}{3(1+2x)}$

The region bounded by the lines $x = -\frac{1}{4}$, $x = \frac{1}{2}$, the x -axis, and the curve is shaded in the figure. This region is rotated through 360 degrees about the x -axis.

- (a) Use calculus to find the exact value of the volume of the solid generated. (5)

Here we shows a paperweight with axis of symmetry AB where $AB = 3$ cm. A is a point on top surface of the paperweight and B is a point on bottom surface of the paperweight. The paperweight is geometrically similar to the solid in part (a).

- (b) Find the volume of this paperweight. (2)

9.

$$I = \int_0^5 e^{\sqrt{3x+1}} \, dx.$$

- (a) Use the substitution $t = \sqrt{3x+1}$ to show that I may be expressed as (5)

$$\int_a^b kte^t \, dx,$$

giving the values of a , b , k .

- (b) Use integration by parts to evaluate this integral, and hence find the value of I correct to 4 significant figures, showing all the steps in your working. (5)

10. Use the substitution $u = 2^x$ to find the exact value of (6)

$$\int_0^1 \frac{2^x}{(2^x + 1)^2} dx.$$

11. (a) Find $\int x \cos 2x dx$. (4)

- (b) Hence, using the identity $\cos 2x = 2 \cos^2 x - 1$, deduce $\int x \cos^2 x dx$. (3)

- 12.

$$\frac{2(4x^2 + 1)}{(2x + 1)(2x - 1)} \equiv A + \frac{B}{2x + 1} + \frac{C}{2x - 1}.$$

- (a) Find the values of A , B , and C . (4)

- (b) Hence show that the exact value of $\int_1^2 \frac{2(4x^2 + 1)}{(2x + 1)(2x - 1)} dx$ is $2 + \ln k$, giving the value of the constant k . (6)

13. Figure 5 shows part of the curve with equation $y = \sqrt{\tan x}$. (4)




Figure 5: $y = \sqrt{\tan x}$

The region bounded R by the lines $x = \frac{\pi}{4}$, the x -axis, and the curve is shaded in the figure. This region is rotated through 2π radians about the x -axis to generate a solid of revolution. Use integration to find an exact value the volume of the solid generated.

14. The curve shown in Figure 6 has equation $y = \frac{1}{2x + 1}$.




Figure 6: $y = \frac{1}{2x + 1}$

The region bounded R by the lines $x = a$, $x = b$, the x -axis, and the curve is shaded in the figure. This region is rotated through 360 degrees about the x -axis to generate a solid of revolution. Find the volume of the solid generated. Express your answer as a single simplified fraction, in terms of a and b .

15. (a) Find $\int \ln\left(\frac{x}{2}\right) dx$. (4)

(b) Find the exact value of $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \sin^2 x dx$. (5)

16. The curve C shown in Figure 7 has parametric equation

$$x = \ln(t + 2), y = \frac{1}{t + 1}, t > -1.$$




Figure 7: $x = \ln(t + 2), y = \frac{1}{t + 1}$

The region bounded R by the lines $x = \ln 2$, $x = \ln 4$, the x -axis, and the curve is shaded in the figure.

(a) Show that the region of R is given by the integral (4)

$$\int_0^2 \frac{1}{(t + 1)(t + 2)} dt.$$

(b) Hence find an exact value for this area. (4)

17. Using the substitution $h = (20 - x)^2$, or otherwise, find the exact value of (6)

$$\int_0^{100} \frac{50}{20 - \sqrt{h}} dh.$$

18. (a) Use integration by parts to find $\int xe^x dx$. (3)

(b) Hence find $\int x^2 e^x dx$. (3)

19. Find $\int_{\frac{\pi}{3}}^{\frac{\pi}{2}} 64 \sin^2 t \cos t dt$, giving your answer in the form $a + b\sqrt{3}$, where a and b are constants to be determined. (4)

20. Figure 8 shows part of the curve $y = \frac{3}{\sqrt{1+4x}}$.




Figure 8: $y = \frac{3}{\sqrt{1+4x}}$

The region bounded R by the lines $x = 0$, $x = 2$, the x -axis, and the curve is shaded in the figure.

(a) Use integration to find the area of R . (4)

The region R is rotated 360° about the x -axis.

(b) Use integration to find the exact value of the volume of the solid formed. (5)

21. (a) Find $\int \tan^2 x dx$. (2)

(b) Use integration by parts to find $\int \frac{1}{x^3} \ln x dx$. (4)

(c) Use the substitution $u = 1 + e^x$ to show that (7)

$$\int \frac{e^{3x}}{1 + e^x} dx = \frac{1}{2}e^{2x} - e^x + \ln(1 + e^x) + k,$$

where k is a constant.

22. Use integration to find the exact area of $y = 3 \cos(\frac{x}{3})$, $0 \leq x \leq \frac{3\pi}{2}$. (3)

23.

$$f(x) = \frac{4 - 2x}{(2x + 1)(x + 1)(x + 3)} = \frac{A}{2x + 1} + \frac{B}{x + 1} + \frac{C}{x + 3}.$$

(a) Find the values of the constants A , B , and C . (4)

(b) Hence find $\int f(x) dx$. (3)

(c) Find $\int_0^2 f(x) dx$ in the form $\ln k$, where k is a constant. (3)

24. (a) Find $\int \sqrt{5-x} dx$. (2)

Figure 9 shows a sketch of the curve with equation

$$y = (x - 1)\sqrt{5 - x}, \quad 1 \leq x \leq 5.$$




Figure 9: $y = (x - 1)\sqrt{5 - x}$

(b) Using integration by parts, or otherwise, find (4)

$$\int (x - 1)\sqrt{5 - x} dx.$$

(c) Hence find (2)

$$\int_1^5 (x - 1)\sqrt{5 - x} dx.$$

25. (a) Using the identity $\cos 2\theta \equiv 1 - 2\sin^2 \theta$, find $\int \sin^2 \theta d\theta$. (2)

Figure 10 shows part of the curve C with parametric equations

$$x = \tan \theta, y = 2 \sin 2\theta, 0 \leq \theta < \frac{\pi}{2}.$$




Figure 10: $x = \tan \theta, y = 2 \sin 2\theta$

The region bounded R by the lines $x = \frac{1}{\sqrt{3}}$, the x -axis, and the curve is shaded in the figure. This region is rotated through 2π radians about the x -axis to generate a solid of revolution.

(b) Show that the volume of the solid of revolution formed is given by the integral (5)

$$k \int_0^{\frac{\pi}{6}} \sin^2 \theta \, d\theta,$$

where k is a constant.

(c) Hence find the exact value for this volume, giving your answer in the form $p\pi^2 + q\pi\sqrt{3}$, where p and q are constants. (3)

26.

$$y = x \ln x, 1 \leq x \leq 4.$$

(a) Use integration by parts to find $\int x \ln x \, dx$. (4)

(b) Hence find the exact area of this integral, giving your answer in the form $\frac{1}{4}(a \ln 2 + b)$, where a and b are integers. (3)

27. Find $\int \frac{9x + 6}{x} \, dx, x > 0$. (2)

28. Using the substitution $x = 2 \cos u$, or otherwise, find the exact value of (7)

$$\int_1^{\sqrt{2}} \frac{1}{x^2 \sqrt{4 - x^2}} \, dx.$$

29. Using the substitution $u = \cos x + 1$, or otherwise, show that

$$\int_0^{\frac{\pi}{2}} e^{\cos x + 1} \sin x \, dx = e(e - 1).$$

30.

$$f(\theta) = 4 \cos^2 \theta - 3 \sin^2 \theta.$$

(a) Show that

$$f(\theta) = \frac{1}{2} + \frac{7}{2} \cos 2\theta. \quad (3)$$

(b) Hence, using calculus, find the exact value of

$$\int_0^{\frac{\pi}{2}} \theta f(\theta) \, d\theta. \quad (7)$$

31. Use integration to find the exact value of

$$\int_0^{\frac{\pi}{2}} x \sin 2x \, dx.$$

32. (a) Express $\frac{5}{(x-1)(3x+2)}$ in partial fractions. (3)

(b) Hence find $\int \frac{5}{(x-1)(3x+2)} \, dx$, where $x > 1$. (3)

33. The curve C has parametric equations

$$x = \ln t, \quad y = t^2 - 2, \quad t > 0.$$




Figure 11: $x = \ln t, y = t^2 - 2$

The region bounded R by the lines $x = \ln 2$, $x = \ln 4$, the x -axis, and the curve is shaded in the figure. This region is rotated through 360° about the x -axis to generate a solid of revolution. Use calculus to find the exact volume of the solid generated.

34. Using the substitution $x = (u - 1)^2 + 1$, or otherwise, and integrating, find the exact value of

$$\int_2^5 \frac{1}{4 + \sqrt{x-1}} dx.$$

35. Figure 12 shows a sketch of the curve with equation $y = x^3 \ln(x^2 + 2)$, $x \geq 0$.




Figure 12: $y = x^3 \ln(x^2 + 2)$

The finite region R , shown shaded in the figure, is bounded by the curve, the x -axis, and the line $x = \sqrt{2}$.

- (a) Use the substitution $u = x^2 + 2$ to show that the area of R is (4)

$$\frac{1}{2} \int_2^4 (u - 2) \ln 2 \, du.$$

- (b) Hence, or otherwise, find the exact area of R . (6)

36. Figure 13 shows part of the curve C with parametric equations

$$x = \tan t, \quad y = \sin t, \quad 0 \leq t < \frac{\pi}{2}.$$




Figure 13: $x = \tan t$, $y = \sin t$

The point P lies on C and has coordinates $(\sqrt{3}, \frac{1}{3}\sqrt{3})$.

- (a) Find the value of t at the point P . (2)

The region bounded R by the lines $x = \sqrt{3}$, the x -axis, and the curve is shaded in the figure. This region is rotated through 2π radians about the x -axis to generate a solid of revolution.

- (b) Find the exact value for the solid of revolution, giving your answer in the form $p\pi\sqrt{3} + q\pi^2$, where p and q are constants. (7)

37. Find $\int (4y + 3)^{-\frac{1}{2}} dy$. (2)

38. (a) Use integration by parts to find $\int x \sin 3x dx$. (3)

(b) Using your answer to part (a), find $\int x^2 \cos 3x dx$. (4)

39. Figure 14 shows the curve with equation (5)

$$y = \sqrt{\frac{2x}{3x^2 + 4}}, \quad x \geq 0.$$




Figure 14: $y = \sqrt{\frac{2x}{3x^2 + 4}}$

The region bounded R by the lines $x = 2$, the x -axis, and the curve is shaded in the figure. This region is rotated through 360° about the x -axis to generate a solid of revolution. Find the exact value for the solid of revolution, giving your answer in the form $k \ln a$, where k and a are constants.

40. Use the substitution $u = 1 + \cos x$, or otherwise, show that (5)

$$\int \frac{2 \sin 2x}{1 + \cos x} dx = 4 \ln(1 + \cos x) - 4 \cos x + k,$$

where k is a constant.

41.

$$f(x) = \frac{1}{x(3x-1)^2} = \frac{A}{x} + \frac{B}{3x-1} + \frac{C}{(3x-1)^2}.$$

(a) Find the value of the constants A , B , and C . (4)

(b) Hence find $\int f(x) dx$. (3)

(c) Find $\int_1^2 f(x) dx$, leaving your answer in the form $a + \ln b$, where a and b are constants. (3)

42. Figure 15 shows a sketch of part of the curve with equation $y = x^{\frac{1}{2}} \ln 2x$.




Figure 15: $y = x^{\frac{1}{2}} \ln 2x$

The finite region R , shown shaded in the figure, is bounded by the curve, the x -axis, and the lines $x = 1$ and $x = 4$.

(a) Find $\int x^{\frac{1}{2}} \ln 2x dx$. (4)

(b) Hence find the exact area of R , giving your answer in the form $a \ln 2 + b$, where a and b are exact constants. (3)

43. (a) Use integration to find (5)

$$\int \frac{1}{x^3} \ln x dx.$$

(b) Hence calculate (2)

$$\int_1^2 \frac{1}{x^3} \ln x dx.$$

44. Figure 16 shows a sketch of part of the curve with equation $y = \frac{x}{1 + \sqrt{x}}$. (8)




Figure 16: $y = \frac{x}{1 + \sqrt{x}}$

The finite region R , shown shaded in the figure, is bounded by the curve, the x -axis, and the lines $x = 1$ and $x = 4$. Use the substitution $u = 1 + \sqrt{x}$, to find, by integrating, the exact area of R .

45. Figure 17 shows a sketch of part of the curve with parametric equations

$$x = 1 - \frac{1}{2}t, y = 2^t - 1.$$




Figure 17: $x = 1 - \frac{1}{2}t, y = 2^t - 1$

- (a) Find the x -coordinate of the point A . (2)

The finite region R , shown shaded in the figure, is bounded by the curve C , the x -axis, and the line $x = -1$.

- (b) Use integration to find exact area of R . (6)

46. Figure 18 shows a sketch of part of the curve with equation $y = 1 - 2 \cos x$. The curve crosses the x -axis at A and B .




Figure 18: $y = 1 - 2 \cos x$

- (a) Find, in terms of π , the x -coordinate of the point A and of the point B . (3)

The region bounded R by the lines the x -axis and the curve is shaded in the figure. This region is rotated through 2π radians about the x -axis to generate a solid of revolution.

- (b) Find, by integration, the exact value for the volume of the solid generated. (6)

47. (a) Find $\int x^2 e^x dx$. (5)

- (b) Hence find the exact value of $\int_0^1 x^2 e^x dx$. (2)

48. Figure 19 shows a sketch of part of the curve with equation (4)

$$y = \sec\left(\frac{1}{2}x\right), 0 \leq x \leq \frac{\pi}{2}.$$

The curve crosses the x -axis at A and B .




Figure 19: $y = \sec\left(\frac{1}{2}x\right)$

The region bounded R by the lines the x -axis, the y -axis, the line $x = \frac{\pi}{2}$, and the curve is shaded in the figure. This region is rotated through 2π radians about the x -axis to generate a solid of revolution. Find, by integration, the exact value for the volume of the solid generated.

49. (a) Use the substitution $x = u^2$, $u > 0$, to show that (3)

$$\int \frac{1}{x(2\sqrt{x} - 1)} dx = \int \frac{2}{u(2u - 1)} du.$$

(b) Hence show that

$$\int_1^9 \frac{1}{x(2\sqrt{x}-1)} dx = 2 \ln\left(\frac{a}{b}\right), \quad (7)$$

where a and b are integers to be determined.

50. Using the substitution $u = 2 + \sqrt{2x+1}$, or other suitable substitutions, find the exact value of

$$\int_0^4 \frac{1}{2 + \sqrt{2x+1}} dx,$$

giving your answer in the form $A + B \ln 2$, where A is an integer and B is a positive constant.

51. Figure 20 shows a sketch of part of the curve with equation $x = 4te^{-\frac{1}{3}t} + 3$. The region bounded R by the lines the x -axis, the t -axis, the line $t = 8$, and the curve is shaded in the figure. (6)




Figure 20: $x = 4te^{-\frac{1}{3}t} + 3$

Use calculus to find the exact value for the area of R .

52. Figure 21 shows a sketch of part of the curve C with parametric equations

$$x = 27 \sec^3 t, \quad y = 3 \tan t, \quad 0 \leq t \leq \frac{\pi}{3}.$$




Figure 21: $x = 27 \sec^3 t, y = 3 \tan t$

- (a) Show that the cartesian equation of C may be written in the form (3)

$$t = \left(x^{\frac{2}{3}} - 9\right)^{\frac{1}{2}}, \quad a \leq x \leq b,$$

stating the values of a and b .

The region bounded R by the lines the x -axis, the line $x = 125$, and the curve is shaded in the figure. This region is rotated through 2π radians about the x -axis to generate a solid of revolution.

- (b) Find, by integration, the exact value for the volume of the solid generated. (5)

53. Use the substitution $u = \sqrt{x}$, or otherwise, to find the exact value of

$$\int_1^4 \frac{10}{2x + 5\sqrt{x}} dx.$$

54. (a) Find $\int xe^{4x} dx$. (3)

- (b) Find $\int \frac{8}{(2x-1)^3} dx, x > \frac{1}{2}$. (2)

55. Figure 22 shows a sketch of part of the curve C with parametric equations

$$x = 3 \tan \theta, \quad y = 4 \cos^2 \theta, \quad 0 \leq \theta < \frac{\pi}{2}.$$




Figure 22: $x = 3 \tan \theta, y = 4 \cos^2 \theta$

The point P lies on C and has coordinates $(3, 2)$. The line l is the normal to C at P . The normal cuts the x -axis at point Q .

- (a) Find the x -coordinate of the point Q . (6)

The region bounded R by the lines the x -axis, the line y -axis, the line l , and the curve is shaded in the figure. This region is rotated through 2π radians about the x -axis to generate a solid of revolution.

- (b) Find the exact value for the solid of revolution, giving your answer in the form $p\pi + q\pi^2$, where p and q are rational numbers to be determined. (9)

56. Figure 23 shows a sketch of part of the curve C with equation (5)

$$y = (2 - x)e^{2x}, x \in \mathbb{R}.$$




Figure 23: $y = (2 - x)e^{2x}$

The region bounded R by the lines the x -axis, the line y -axis, and the curve is shaded in the figure. Use calculus, showing each step in your working, to obtain an exact value for the area of R . Give your answer in its simplest form.

57. (a) Express $\frac{25}{x^2(2x + 1)}$ in partial fractions. (4)

Figure 24 shows a sketch of part of the curve C with equation

$$y = \frac{5}{x\sqrt{2x+1}}.$$




Figure 24: $y = \frac{5}{x\sqrt{2x+1}}$

The region bounded R by the lines the x -axis, the line $x = 1$, the line $x = 4$, and the curve is shaded in the figure. This region is rotated through 360° about the x -axis.

- (b) Use calculus to find the exact value for the solid of revolution, giving your answer in the form $a + b \ln c$, where a , b , and c are constants. (6)

58. Figure 25 shows a sketch of part of the curve with equation

$$y = 4x - xe^{\frac{1}{2}x}, \quad x \geq 0.$$




Figure 25: $y = 4x - xe^{\frac{1}{2}x}$

The curve meets the x -axis the the origin O and cuts the x -axis at the point A .

- (a) Find, in terms of $\ln 2$, the x -coordinate of the point A . (2)

(b) Find $\int x e^{\frac{1}{2}x} dx$. (3)

The region bounded R by the lines the x -axis and the curve with equation

$$y = 4x - x e^{\frac{1}{2}x}, \quad x \geq 0.$$

(c) Find, by integration, the exact value for the area of R . Give your answer in terms of $\ln 2$. (3)

59. Figure 26 shows a sketch of part of the curve with equation

$$y = \sqrt{(3-x)(x+1)}, \quad 0 \leq x \leq 3.$$




Figure 26: $y = \sqrt{(3-x)(x+1)}$

The region bounded R by the lines the x -axis, the y -axis, and the curve.

(a) Use the substitution $x = 1 + 2 \sin \theta$ to show that (5)

$$\int_0^3 \sqrt{(3-x)(x+1)} dx = k \int_{-\frac{\pi}{6}}^{\frac{\pi}{2}} \cos^2 \theta d\theta,$$

where k is a constant to be determined.

(b) Hence find, by integration, the exact value of R . (3)

60. Figure 27 shows a sketch of part of the curve with equation

$$y = 3^x.$$




Figure 27: $y = 3^x$

The point P lies on C and has coordinates $(2, 9)$. The line l is a tangent to C at P . The line l cuts the x -axis at the Q .

- (a) Find the exact value of the x -coordinate of Q . (4)

The region bounded R by the lines the x -axis, the line y -axis, the line l , and the curve is shaded in the figure. This region is rotated through 360° about the x -axis.

- (b) Use integration to find the exact value of the volume of the solid generated. Give your answer in the form $\frac{p}{q}$, where p and q are exact constants. (6)

61. Figure 28 shows a sketch of part of the curve with equation

$$y = x^2 \ln x, \quad x \geq 1.$$




Figure 28: $y = x^2 \ln x$

The region bounded R by the lines the x -axis, the line $x = 2$ -axis, and the curve is shaded in the figure. Use integration to find the exact value of the area of R .

62. (a) Given that $y > 0$, find (6)

$$\int \frac{3y - 4}{y(3y + 2)} dy.$$

- (b) Use the substitution $x = 4 \sin^2 \theta$ to show that (5)

$$\int_0^3 \sqrt{\frac{x}{4-x}} dx = \lambda \int_0^{\frac{\pi}{3}} \sin^2 \theta d\theta,$$

where λ is a constant to be determined.

- (c) Hence use integration to find (4)

$$\int_0^3 \sqrt{\frac{x}{4-x}} dx,$$

giving your answer in the form $a\pi + b$, where a and b are exact constants.

63. (a) Find (2)

$$\int (2x - 1)^{\frac{3}{2}} dx,$$

giving your answer in its simplest form.

Figure 29 shows a sketch of part of the curve with equation

$$y = (2x - 1)^{\frac{3}{4}}, \quad x \geq \frac{1}{2}.$$




Figure 29: $(2x - 1)^{\frac{3}{4}}$

The curve C cuts the line $y = 8$ at the point P with coordinates $(k, 8)$, where k is a constant.

- (b) Find the value of k . (2)

The region bounded R by the lines the x -axis, the line y -axis, the line $y = 8$, and the curve is shaded in the figure. This region is rotated through 2π radians about the x -axis.

- (c) Use integration to find the exact value of the volume of the solid generated. (4)

64. Figure 30 shows a sketch of part of the curve with equation

$$y = \frac{6}{e^x + 2}, \quad x \in \mathbb{R}.$$




Figure 30: $y = \frac{6}{e^x + 2}$

The finite region R , shown shaded in Figure 30, is bounded by the curve, the y -axis, the x -axis, and the line with equation $x = 1$.

(a) Use the substitution $u = e^x$ to show that the area of R can be given by (2)

$$\int_a^b \frac{6}{u(u+2)} du,$$

where a and b are constants to be determined.

(b) Hence use calculus to find the exact area of R . (6)

65. The finite region S , shown shaded in Figure 31, is bounded by the y -axis, the x -axis, the line with equation $x = \ln 4$, and the curve with equation (7)

$$y = e^x + 2e^{-x}, \quad x \geq 0.$$




Figure 31: $y = e^x + 2e^{-x}$

The region S is rotated through 2π radians about the x -axis. Use integration to find the exact value of the volume of the solid generated. Give your answer in its simplest form.

66. Figure 32 shows a sketch of part of the curve C with parametric equations

$$x = 3\theta \sin \theta, y = \sec^3 \theta, 0 \leq \theta < \frac{\pi}{2}.$$




Figure 32: $x = 3\theta \sin \theta, y = \sec^3 \theta$

The point $P(k, 8)$ lies on C , where k is a constant.

(a) Find the exact value of k . (2)

The finite region R , shown shaded in Figure 32, is bounded by the curve C , the y -axis, the x -axis, and the line with equation $x = k$.

(b) Show that the area of R can be expressed in the form (4)

$$\lambda \int_{\alpha}^{\beta} (\theta \sec^2 \theta + \tan \theta \sec^2 \theta) d\theta,$$

where α , β , and λ are constants to be determined.

(c) Hence use integration to find the exact value of the area of R . (6)

67. Figure 33 shows a vertical cylindrical tank of height 200 cm containing water. Water is leaking from a hole P on the side of the tank.




Figure 33: a leaky tank

At time t minutes after the leaking starts, the height of water in the tank is h cm. The height h cm of the water in the tank satisfies the differential equation

$$\frac{dh}{dt} = k(h - 9)^{\frac{1}{2}}, \quad 9 < h \leq 200,$$

where k is a constant. Given that, when $h = 130$, the height of the water is falling at a rate of 1.1 cm per minute,

- (a) find the value of k . (2)

Given that the tank was full of water when the leaking started,

- (b) solve the differential equation with your value of k , to find the value of t when $h = 50$. (6)