

Dr Oliver Mathematics
Cambridge O Level Additional Mathematics
2010 November Paper 1 Variant 3: Calculator
2 hours

The total number of marks available is 80.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

You must write down all the stages in your working.

1. Show that (3)

$$\sec x - \cos x \equiv \sin x \tan x.$$

2. A 4-digit number is formed by using four of the seven digits 2, 3, 4, 5, 6, 7, and 8.
No digit can be used more than once in any one number.

Find how many different 4-digit numbers can be formed if

- (a) there are no restrictions, (2)

- (b) the number is even. (2)

3. The line (4)

$$y = mx + 2$$

is a tangent to the curve

$$y = x^2 + 12x + 18.$$

Find the possible values of m .

4. The remainder when the expression (4)

$$x^3 + kx^2 - 5x - 3$$

is divided by $(x - 2)$ is 5 times the remainder when the expression is divided by $(x + 1)$.

Find the value of k .

5. Solve the simultaneous equations (5)

$$\log_3 a = 2 \log_3 b$$

$$\log_3(2a - b) = 1.$$

6. Solve the equation (6)

$$3x^3 + 7x^2 - 22x - 8 = 0.$$

7. (a) Sketch the graph of (3)

$$y = |3x - 5| \text{ for } -2 \leq x \leq 3,$$

showing the coordinates of the points where the graph meets the axes.

- (b) On the same diagram, sketch the graph of (1)

$$y = 8x.$$

- (c) Solve the equation (3)

$$8x = |3x - 5|.$$

8. A function f is defined, for $x \in \mathbb{R}$, by

$$f(x) = x^2 + 4x - 6.$$

- (a) (i) Find the least value of $f(x)$ and the value of x for which it occurs. (2)

- (ii) Hence write down a suitable domain for $f(x)$ in order that $f^{-1}(x)$ exists. (1)

- (b) Functions g and h are defined, for $x \in \mathbb{R}$, by

$$g(x) = \frac{1}{2}x - 1,$$

$$h(x) = x^2 - x.$$

- (i) Find $g^{-1}(x)$. (2)

- (ii) Solve (3)

$$g h(x) = g^{-1}(x).$$

9. (a) Find (3)

$$\int \left(x^{\frac{1}{3}} - 3 \right)^2 dx.$$

- (b) (i) Given that (3)

$$y = x\sqrt{x^2 + 6},$$

find $\frac{dy}{dx}$.

- (ii) Hence find (2)

$$\int \frac{x^2 + 3}{\sqrt{x^2 + 6}} dx.$$

10. A particle travels in a straight line so that, t s after passing through a fixed point O , its displacement s m from O is given by

$$s = \ln(t^2 + 1).$$

- (a) Find the value of t when $s = 5$. (2)
- (b) Find the distance travelled by the particle during the third second. (2)
- (c) Show that, when $t = 2$, the velocity of the particle is 0.8 ms^{-1} . (2)
- (d) Find the acceleration of the particle when $t = 2$. (3)

11. Solve the equation

- (a) $3 \sin x - 4 \cos x = 0$, for $0^\circ \leq x \leq 360^\circ$, (3)
- (b) $11 \sin y + 1 = 4 \cos^2 y$, for $0^\circ \leq y \leq 360^\circ$, (4)
- (c) $\sec(2z + \frac{1}{3}\pi) = -2$, for $0 \leq z \leq \pi$ radians. (4)

EITHER

12. A curve has the equation

$$y = A \sin 2x + B \cos 3x.$$


The curve passes through the point with coordinates $(\frac{1}{12}\pi, 3)$ and has a gradient of -4 when $x = \frac{1}{3}\pi$.

- (a) Show that $A = 4$ and find the value of B . (6)
- (b) Given that, for $0 \leq x \leq \frac{1}{3}\pi$, the curve lies above the x -axis, find the area of the region enclosed by the curve, the y -axis, and the line $x = \frac{1}{3}\pi$. (5)

OR

13. The diagram shows the curve

$$y = 4x^2 - 2x^3.$$



- The point A lies on the curve and the x -coordinate of A is 1.
 - The curve crosses the x -axis at the point B .
 - The normal to the curve at the point A crosses the y -axis at the point C .
- (a) Show that the coordinates of C are $(0, 2.5)$. (5)
- (b) Find the area of the shaded region. (6)

*Dr Oliver
Mathematics*

*Dr Oliver
Mathematics*

*Dr Oliver
Mathematics*

*Dr Oliver
Mathematics*

*Dr Oliver
Mathematics*